

# 海桐花属一些种类的新异名

<sup>1</sup> 张志耘 <sup>2</sup> Nicholas J. Turland

<sup>1</sup>(中国科学院植物研究所系统与进化植物学开放研究实验室 北京 100093)

<sup>2</sup>(Missouri Botanical Garden, P. O. Box 299, St. Louis, Missouri 63166-0299, U.S. A.)

## New synonymies of some species of *Pittosporum* (Pittosporaceae)

<sup>1</sup>ZHANG Zhi-Yun <sup>2</sup>Nicholas J. Turland

<sup>1</sup>(Laboratory of Systematic and Evolutionary Botany, the Chinese Academy of Sciences, Beijing 100093)

<sup>2</sup>(Missouri Botanical Garden, P. O. Box 299, St. Louis, Missouri 63166-0299, U.S. A.)

**Abstract** As part of a revision of Chinese *Pittosporum* Banks ex Gaertn. for the forthcoming account of Pittosporaceae in "Flora of China", Volume 9, eight names, all described from China, are reduced to synonymy as follows: *P. illicioides* Makino var. *angustifolium* T. C. Huang ex S. Y. Lu, syn. nov. and *P. illicioides* var. *stenophyllum* P. L. Chiu ex H. T. Chang & S. Z. Yan, syn. nov., both under *P. illicioides*; *P. densinervatum* H. T. Chang & S. Z. Yan, syn. nov. and *P. longicarpum* S. K. Wu ex C. Y. Wu, syn. nov., both under *P. kweichowense* Gowda var. *kweichowense*; *P. polycarpum* H. T. Chang & S. Z. Yan, syn. nov. under *P. paniculiferum* H. T. Chang & S. Z. Yan; *P. membranifolium* S. C. Huang ex C. Y. Wu, syn. nov. under *P. perryanum* Gowda var. *perryanum*; *P. monanthum* C. Y. Wu, syn. nov. under *P. podocarpum* Gagnep. var. *podocarpum*; and *P. tobira* (Thunb.) W. T. Aiton var. *chinense* S. Kobayashi, syn. nov. under *P. tobira*.

**Key words** Pittosporaceae; *Pittosporum*; New synonymies

### 1 海金子

***Pittosporum illicioides*** Makino in Bot. Mag. Tokyo 14: 32. 1900. TYPE: Japan. Harima: Kishima-mura in Ibo-gori, Feb 1900, U. Ouye s.n. (holotype, not seen).

*P. illicioides* var. *angustifolium* T. C. Huang ex S. Y. Lu in Quart. J. Chin. Forest. 10: 144. 1977, syn. nov. TYPE: China. Taiwan (台湾), Taizhong (台中), Heping Xiang, Bi Lu Xi, without date (fr.), S. Y. Lǚ (吕胜由) 5916 (holotype, TAIF, not seen).

*P. illicioides* var. *stenophyllum* P. L. Chiu ex H. T. Chang & S. Z. Yan in Fl. Reip. Pop. Sin. 35(2): 16. 1979, syn. nov. TYPE: China. Zhejiang (浙江), Longquan (龙泉), streamside in valley, 700 m, without date, S. Y. Zhang (章绍尧) 3262 (holotype, HHBG, not seen).

*P. illicioides* var. *stenophyllum* is considered to differ from *P. illicioides* var. *illicioides* in having leaves narrowly lanceolate, 10~18 cm × 1.7~3.3 cm. However, these characters fall with-

in the range of variation of typical *P. illicioides*. *P. illicioides* var. *angustifolium* (described from Taiwan) represents the extreme of the continuous range of variation, with leaves linear, 5 ~ 15 cm  $\times$  0.6 ~ 2 cm.

Distribution: China (Anhui, Fujian, Guangdong, Guangxi, Guizhou, Hubei, Hunan, Jiangsu, Jiangxi, Sichuan, Taiwan, and Zhejiang), Japan.

In forests, thickets, valleys, and along streamsides from 100 ~ 2200 m in altitude.

## 2 贵州海桐

***Pittosporum kweichowense*** Gowda, J. Arnold Arbor. 32: 296. 1951. TYPE: China. Guizhou (贵州), Anlong "Anlung" (安龙), 500 m, light woods, 1930-10-20, Y. Tsiang (蒋英) 7444 (holotype, NY, not seen).

*P. densinervatum* H. T. Chang & S. Z. Yan in Acta Phytotax. Sin. 16(4): 86. 1978, syn. nov. TYPE: China. Hunan (湖南), Longshan (龙山), forest margin, 1060 m, without date, L. H. Liu (刘林翰) 1942 (holotype, IBSC!).

*P. longicarpum* S. K. Wu ex C. Y. Wu, Fl. Yunnan. 3: 314, pl. 90: 2. 1983, syn. nov. TYPE: China. Yunnan (云南): Wenshan (文山), forest, 1958-09-24 (fr.), H. T. Tsai (蔡希陶) 58-8053 (holotype, KUN; photo, PE!).

The original description of *P. densinervatum* indicated that it differs from *P. kweichowense* in having elliptic leaves 2 ~ 3 cm wide. However, this character is quantitative and clearly falls within the range of variation of *P. kweichowense*. *P. longicarpum* is considered to differ from *P. kweichowense* in having oblong capsules 15 ~ 20 mm long, and longer fruiting pedicels. However, based on our studies these characters show continuous variation and there are no other characters that distinguish *P. kweichowense* from *P. longicarpum* S. K. Wu ex C. Y. Wu.

Distribution: China [SW Guizhou (Anlong), W Hunan (Zhijiang, Longshan), and SE Yunnan (Wenshan)].

In forests, thickets, and along riversides from 500 ~ 2000 m in altitude.

## 3 圆锥海桐

***Pittosporum paniculiferum*** H. T. Chang & S. Z. Yan in Acta Phytotax. Sin. 16(4): 90. 1978. TYPE: China. Yunnan (云南), Lincang (临沧), Mayidui, without date, J. S. Xin (辛景三) 406 (syntype, IBSC!); *ibid.*, J. S. Xin (辛景三) 523 (syntype, KUN, not seen).

*P. polycarpum* H. T. Chang & S. Z. Yan, Acta Phytotax. Sin. 16(4): 90. 1978, syn. nov. TYPE: China. Sichuan (四川), Mt. Emei (峨眉山), without date, Y. X. Xiao (肖永贤) 49736 (syntype, SYS!); *ibid.*, G. H. Yang (杨光辉) 54650 (syntype, SZ, not seen).

*P. polycarpum* is considered to differ from *P. paniculiferum* in having capsules subglobose and seeds more numerous; however, these character states also occur in the latter species. *P. polycarpum* was omitted from the treatment of the genus in Flora Reipublicae Popularis Sinicae (Chang & Yan, 1979).

Distribution: China [C Sichuan (Mt. Emei) and S Yunnan].

In forests, thickets, and on limestone slopes from 500 ~ 1600 m in altitude.

## 4 缝线海桐

***Pittosporum perryanum*** Gowda in J. Arn. Arb. 32: 290. 1951. TYPE: China. Guangxi

(广西), Shiwandashan (Sup Man Ta Shan) (十万大山), 1937-07-21, H. Y. Liang (梁向日) 69823 (holotype, A, not seen, isotype, IBSC!).

*P. membranifolium* S. C. Huang ex C. Y. Wu, Fl. Yunnan. 3: 314. Pl. 90: 3. 1983, syn. nov. TYPE: China. Yunnan (云南), Shuangjiang (双江), 1957-09-20 (fr.), J. S. Xin (辛景三) 1089 (holotype, KUN; photo, PE!).

*P. membranifolium* is considered to differ from *P. perryanum* in having leaves longer, 16 ~ 22 cm, apex cuspidate to abruptly acuminate, and capsules larger, 25 ~ 32 mm × 14 ~ 17 mm. However, these morphological characters also occur in *P. perryanum*, and thus the establishment of *P. membranifolium* is not supported.

Distribution: China [W Guangdong (Xinyi), S Guangxi, SW Hainan, S Sichuan (Changning, Hejiang), and S Yunnan].

In forests, thickets, valleys, and on slopes and stream sides from 600 ~ 1800 m in altitude.

## 5 柄果海桐

*Pittosporum podocarpum* Gagnep. in Notul. Syst. (Paris) 8: 211. 1939. TYPE: China. W. China (中国西部), 1904, Wilson 3231 (syntype, P, not seen); ibid., Wilson 3231a (syntype, P, not seen); Chongqing (重庆), Chengkou "district Tchen-keou-tin" (城口), [Chengkou], without date (fl. & fr.), Farges 1166 (syntype, P, not seen); Yunnan (云南), Binchuan (宾川) "Pin-tchouan" [Binchuan], without date, Ducloux 6709 (syntype, P, not seen).

*P. monanthum* C. Y. Wu, Fl. Yunnan. 3: 318. Pl. 91: 2. 1983, syn. nov. TYPE: China. Yunnan (云南), Jingdong (景东), forest, 1963-05-15 (fl.), C. A. Wu (武全安) 9222 (holotype, KUN; photo, PE!).

The original description of *P. monanthum* indicated that it is close to *P. podocarpum* but differs in having leaves oblanceolate, lateral veins few, in 4 or 5 pairs, flowers solitary, pedicels very shortly pilose, and calyx cupular and pilose. We have observed all these characters in several specimens of *P. podocarpum* from Sichuan, and found that these characters are quantitative and clearly fall within the variation range of *P. podocarpum*.

Distribution: China (Fujian, Gansu, Guangxi, Guizhou, Hubei, Hunan, Shaanxi, Sichuan, Xizang, and Yunnan), India, Myanmar and Vietnam.

In forests and shady places, and on slopes and riversides from 500 ~ 2700 (~ 3000) m in altitude.

## 6 海桐

*Pittosporum tobira* (Thunb.) W. T Aiton, Hort. Kew. 2: 27. 1811. — *Euonymus tobira* Thunb. in Nova Acta Regiae Soc. Sci. Upsal. 3: 208. 1780. TYPE: described from Japan, not seen.

*P. tobira* var. *chinense* S. Kobayashi in J. Jpn. Bot. 57: 74. 1982, syn. nov. TYPE: China. Jiangsu (江苏), Suzhou (苏州), without date, S. Oka 359 (holotype, TI, not seen).

The original description of *P. tobira* var. *chinense* indicated that it is close to *P. tobira* var. *tobira*, but in the former inflorescences having appressed, short, soft hairs, while in the latter inflorescences having long-armed hairs, and the stalks are composed of 5 ~ 7 cells. Except the type specimen collected from Suzhou of Jiangsu Province, up to now there is no record from China. How-

ever, in the specimens of Kyushu and Korea, short-and long-armed hairs are mixed about half and half on an inflorescence (Kobayashi, 1982). Therefore, we are reluctant to accept var. *chinense* as a distinct taxon, and prefer to regard it as a synonym of *P. tobira* var. *tobira*.

Distribution: China (Fujian, Guangdong, Guangxi, Guizhou, Hainan, Hubei, Jiangsu, Sichuan, Yunnan, and Zhejiang).

**Acknowledgments** We would like to thank Anthony Brach (Harvard University Herbaria), Deng Yun-Fei (South China Institute of Botany, the Chinese Academy of Sciences), and Hideaki Ohba (University of Tokyo) for checking type specimens at their respective Herbaria. We also thank Peng Hua (Kunming Institute of Botany, the Chinese Academy of Sciences), Yang Qin-Er (Institute of Botany, the Chinese Academy of Sciences) and Zhu Guang-Hua (Missouri Botanical Garden, St. Louis, USA) for advice and help.

### References

- Chang H-T (张宏达), Yan S-Z (颜素珠), 1979. Hamamelicaceae. In: Flora Reipublicae Popularis Sinicae. Beijing: Science Press. 35(2): 1 ~ 36
- Kobayashi Sumiko, 1982. A taxonomical note on *Pittosporum tobira* and its allied species. J Jpn Bot, 57: 70 ~ 80

**摘要** 作为对英文版“Flora of China”海桐花科的分类学修订结果,提出了海桐花属 6 个种的 8 个新异名,并分别进行了讨论。

**关键词** 海桐花科;海桐花属;新异名

(责任编辑 白羽红)